



Original Effective Date: 04/2012  
Current Effective Date: 02/25/2023  
Last P&T Approval/Version: 01/31/2024  
Next Review Due By: 01/2025  
Policy Number: C4719-A

## Serotonin - Norepinephrine Reuptake Inhibitors (SNRI)

### PRODUCTS AFFECTED

desvenlafaxine tab ER, FETZIMA (levomilnacipran), KHEDEZLA (desvenlafaxine), PRISTIQ (desvenlafaxine)

### COVERAGE POLICY

*Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Coverage Guideline must be read in its entirety to determine coverage eligibility, if any. This Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide Molina Healthcare complete medical rationale when requesting any exceptions to these guidelines.*

#### **Documentation Requirements:**

*Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.*

#### **DIAGNOSIS:**

Treatment of Major Depressive Disorder in Adults, Treatment of Hot Flashes associated with Menopause

#### **REQUIRED MEDICAL INFORMATION:**

This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. If a drug within this policy receives an updated FDA label within the last 180 days, medical necessity for the member will be reviewed using the updated FDA label information along with state and federal requirements, benefit being administered and formulary preferencing. Coverage will be determined on a case-by case basis until the criteria can be updated through Molina Healthcare, Inc. clinical governance. Additional information may be required on a case-by-case basis to allow for adequate review. When the requested drug product for coverage is dosed by weight, body surface area or other member specific measurement, this data element is required as part of the medical necessity review. The Pharmacy and Therapeutics Committee has determined that the drug benefit shall be a mandatory generic and that generic drugs will be dispensed whenever available.

#### **A. FOR ALL INDICATIONS:**

1. Product being requested has an FDA labeled indication or compendia supported use for diagnosis, age, and dose  
AND

## Drug and Biologic Coverage Criteria

2. Documentation of an adequate trial (3 months) and therapeutic failure to at least 3 formulary preferred SNRI products.

### CONTINUATION OF THERAPY:

#### A. FOR ALL INDICATIONS:

1. Adherence to therapy at least 85% of the time as verified by the prescriber or member medication fill history  
AND
2. Prescriber attests to or clinical reviewer has found no evidence of intolerable adverse effects or drug toxicity  
AND
3. Documentation of positive clinical response as demonstrated by low disease activity and/or improvements in the condition's signs and symptoms

### DURATION OF APPROVAL:

Initial authorization: 12 months, Continuation of Therapy: 12 months

### PRESCRIBER REQUIREMENTS:

None

### AGE RESTRICTIONS:

18 years of age and older

### QUANTITY:

30 tablets per 30 days

### PLACE OF ADMINISTRATION:

The recommendation is that oral medications in this policy will be for pharmacy benefit coverage and patient self-administered.

## DRUG INFORMATION

### ROUTE OF ADMINISTRATION:

Oral

### DRUG CLASS:

Serotonin – Norepinephrine reuptake inhibitors (SNRI)

### FDA-APPROVED USES:

Indicated for the treatment of Major Depressive Disorder in adults

*Limitations of Use (Fetzima only): Fetzima is not approved for the management of fibromyalgia. The efficacy and safety of Fetzima for the management of fibromyalgia have not been established.*

### COMPENDIAL APPROVED OFF-LABELED USES:

Treatment of Hot Flashes associated with Menopause (Pristiq and Khedezla)

## APPENDIX

### APPENDIX:

**Reserved for State specific information.** Information includes, but is not limited to, State contract language, Medicaid criteria and other mandated criteria.

Molina Healthcare, Inc. confidential and proprietary © 2024

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

### **State Specific Information**

#### **State Marketplace**

**Nevada** (Source: Nevada Legislature)

“Chapter 689A of Nevada Revised Statutes (NRS) is hereby amended by adding thereto a new section to read as follows:

1. A policy of health insurance which provides coverage for prescription drugs must not require an insured to submit to a step therapy protocol before covering a drug approved by the Food and Drug Administration that is prescribed to treat a psychiatric condition of the insured, if:
  - a. The drug has been approved by the Food and Drug Administration with indications for the psychiatric condition of the insured or the use of the drug to treat that psychiatric condition is otherwise supported by medical or scientific evidence;
  - b. The drug is prescribed by:
    - i. A psychiatrist
    - ii. A physician assistant under the supervision of a psychiatrist;
    - iii. An advanced practice registered nurse who has the psychiatric training and experience prescribed by the State Board of Nursing pursuant to NRS 632.120; or
    - iv. A primary care provider that is providing care to an insured in consultation with a practitioner listed in subparagraph (1), (2) or (3), if the closest practitioner listed in subparagraph (1), (2) or (3) who participates in the network plan of the insurer is located 60 miles or more from the residence of the insured; and
  - c. The practitioner listed in paragraph (b) who prescribed the drug knows, based on the medical history of the insured, or reasonably expects each alternative drug that is required to be used earlier in the step therapy protocol to be ineffective at treating the psychiatric condition...
3. As used in this section:
  - c. *‘Step therapy protocol’ means a procedure that requires an insured to use a prescription drug or sequence of prescription drugs other than a drug that a practitioner recommends for treatment of a psychiatric condition of the insured before his or her policy of health insurance provides coverage for the recommended drug.’*

Molina Reviewer Note: Medical necessity review for a psychiatric condition cannot require trial of other medications first. This is applicable to formulary medications that require prior authorization and non-formulary medications and is not limited to only medications designated ‘ST’. If the requested drug is a brand name and the generic is on formulary, request can be reviewed for specific medical reason generic cannot be used.

## BACKGROUND AND OTHER CONSIDERATIONS

### **BACKGROUND:**

#### *Genetic Testing and Psychiatric Medication Choice*

Pharmacogenomics is the study of the relationship between a genetic variation and how the body responds to medication. The American Psychiatric Association (APA) has a statement on pharmacogenomic testing based on results of their own task force looking at novel biomarkers and treatments. This task force determined “there is not sufficient information to support the widespread use of pharmacogenetic testing in clinical practice.” The 2020 International Society of Psychiatric Genetics (ISPG) guidelines on genetic testing states “Pharmacogenetic testing should be viewed as a decision-support tool to assist in thoughtful implementation of good clinical care. We recommend HLA-A and HLA-B testing prior to use of carbamazepine and oxcarbazepine, in alignment with regulatory agencies and expert groups. Evidence to support widespread use of other pharmacogenetic tests at this time is still inconclusive, but when pharmacogenetic testing results are already available, providers are encouraged to integrate this information into their medication selection and dosing decisions. Genetic information for CYP2C19 and CYP2D6 would likely be most beneficial for individuals who have experienced an inadequate response or adverse reaction to a previous antidepressant or antipsychotic trial.” The Pharmacogenomics Knowledgebase (PharmGKB) has summaries of genotype based *dosing* recommendations only, including the following:

- Aripiprazole: The Royal Dutch Pharmacists Association - Pharmacogenetics Working Group

## Drug and Biologic Coverage Criteria

(DPWG) recommends reducing maximum dose of aripiprazole for patients carrying poor metabolizer alleles of CYP2D6.

- Antidepressants: The French National Network of Pharmacogenetics (Réseau national de pharmacogénétique (RNPgX)) recommends CYP2D6 and CYP2C19 genotyping before initiating an antidepressant treatment, especially in patients with a high risk of toxicity.
- Brexpiprazole: The Royal Dutch Pharmacists Association - Pharmacogenetics Working Group (DPWG) recommends to use half of the standard dose of brexpiprazole for patients carrying poor metabolizer alleles of CYP2D6.
- Duloxetine: There are currently no dosing recommendations for duloxetine based on CYP2D6 genotype.

At this time, there are no guidelines found that support the use of pharmacogenomic testing to make therapeutic treatment decisions based on the results of that testing. These tests may be beneficial to help guide dosing and toxicity concerns of specific agents for those with suboptimal responses or at risk of toxicity.

### **CONTRAINDICATIONS/EXCLUSIONS/DISCONTINUATION:**

All other uses of SNRI's are considered experimental/investigational and therefore, will follow Molina's Off- Label policy.

Contraindications to Fetzima (levomilnacipran) include: Hypersensitivity to levomilnacipran, milnacipran HCl, or any excipient in the Fetzima formulation, use with MAOIs intended to treat psychiatric disorders or within 7 days of stopping treatment with Fetzima, use within 14 days of stopping an MAOI intended to treat psychiatric disorders, and patients being treated with linezolid or intravenous methylene blue.

Contraindications to Khedezla (desvenlafaxine) include: hypersensitivity to desvenlafaxine succinate, venlafaxine hydrochloride or any excipients in the Khedezla Extended-release Tablets formulation, use with MAOIs intended to treat psychiatric disorders or within 7 days of stopping treatment with Khedezla, use within 14 days of stopping an MAOI intended to treat psychiatric disorders, and patients being treated with linezolid or intravenous methylene blue.

Contraindications to Pristiq (desvenlafaxine) include: Hypersensitivity to desvenlafaxine succinate, venlafaxine hydrochloride or any excipients in the Pristiq formulation, use with MAOIs intended to treat psychiatric disorders or within 7 days of stopping treatment with Pristiq, use within 14 days of stopping an MAOI intended to treat psychiatric disorders, and patients being treated with linezolid or intravenous methylene blue.

### **OTHER SPECIAL CONSIDERATIONS:**

Fetzima (levomilnacipran), Khedezla (desvenlafaxine), and Pristiq (desvenlafaxine) have a black box warning for suicidal thought and behaviors.

## **CODING/BILLING INFORMATION**

*Note: 1) This list of codes may not be all-inclusive. 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement*

<b>HCPCS CODE</b>	<b>DESCRIPTION</b>
NA	

### **AVAILABLE DOSAGE FORMS:**

Desvenlafaxine ER TB24 50MG, 100MG

Desvenlafaxine Succinate ER TB24 25MG, 50MG, 100MG

Molina Healthcare, Inc. confidential and proprietary © 2024

*This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.*

## Drug and Biologic Coverage Criteria

Fetzima CP24 20MG, 40MG, 80MG 120MG

Fetzima Titration C4PK 20 & 40MG

Khedezla TB24 50MG, 100MG

Pristiq TB24 25MG, 50MG, 100MG

### REFERENCES

1. Fetzima (levomilnacipran) [prescribing information]. Madison, NJ: Allergan USA Inc; October 2023.
2. Khedezla (desvenlafaxine) [prescribing information]. Morristown, NJ: Pernix Therapeutics LLC; August 2023.
3. Pristiq (desvenlafaxine) [prescribing information]. Philadelphia, PA: Wyeth Pharmaceuticals LLC; August 2023.
4. Bauer M, Pfennig A, Severus E, et al. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of unipolar depressive disorders, part 1: update 2013 on the acute and continuation treatment of unipolar depressive disorders. *World J Biol Psychiatry* 2013; 14:334.
5. Kupfer DJ, Frank E, Phillips ML. Major depressive disorder: new clinical, neurobiological, and treatment perspectives. *Lancet* 2012; 379:1045.
6. Marcus SC, Olfson M. National trends in the treatment for depression from 1998 to 2007. *Arch Gen Psychiatry* 2010; 67:1265.
7. News Author, Author, N., & APA Curated and updated for the community by APA. (2019, February 06). Genetic testing to improve psychiatric medication choice. Retrieved December 9, 2022, from <https://www.psychiatry.org/news-room/apa-blogs/genetic-testing-to-improve-psychiatric-medication>
8. Moran, M. (2018). Task Force on gene testing for antidepressant efficacy concludes tests not yet ready for widespread use. *Psychiatric News*, 53(14). doi:10.1176/appi.pn.2018.pp6b1
9. Zeier, Z., Carpenter, L. L., Kalin, N. H., Rodriguez, C. I., McDonald, W. M., Widge, A. S., & Nemeroff, C. B. (2018). Clinical implementation of Pharmacogenetic decision support tools for antidepressant drug prescribing. *American Journal of Psychiatry*, 175(9), 873-886. doi:10.1176/appi.ajp.2018.17111282
10. International Society of Psychiatric Genetics (ISPG) Genetic testing statement. (n.d.). Retrieved December 9, 2022, from <https://ispg.net/genetic-testing-statement/>
11. Pharmacogenomics Knowledgebase (PharmGKB) Clinical Guideline Annotations. (n.d.). Retrieved December 9, 2022, from <https://www.pharmgkb.org/guidelineAnnotations>

SUMMARY OF REVIEW/REVISIONS	DATE
REVISION- Notable revisions: FDA-Approved Uses Available Dosage Forms References	Q1 2024
REVISION- Notable revisions: Products Affected Diagnosis Required Medical Information Continuation of Therapy Quantity Compendial Approved Off-Labeled Uses Background Contraindications/Exclusions/Discontinuation Other Special Considerations Available Dosage Forms References	Q1 2023
Q2 2022 Established tracking in new format	Historical changes on file