

Why Are Immunizations So Important?

Immunizations are shots that help the body fight diseases. Each shot fights a different disease like chickenpox, measles, or mumps. Kids need shots to fight disease and to be ready for school.

Talk to a doctor about the shots your kid needs to be protected from diseases. **Remember, these shots are covered at no cost to you.**

Here's a look at the shot schedule recommended by the Centers for Disease Control and Prevention (CDC):



Hepatitis B (HepB)



HepB (2nd Dose)



Rotavirus (RV)
Diphtheria, Tetanus,
Acellular Pertussis (DTaP)
Haemophilus Influenza
Type B (Hib)
Pneumococcal
Conjugate (PCV13)
Inactivated
Poliovirus (IPV)



RV (2nd Dose)
DTaP (2nd Dose)
Hib (2nd Dose)
PCV13 (2nd Dose)
IPV (2nd Dose)



HepB (3rd Dose 6-18
months)
RV (3rd Dose)
DTaP (3rd Dose)
Hib (3rd Dose)
PCV13 (3rd Dose)
IPV (3rd Dose 6-18 months)



Hib (4th Dose)
PCV13 (4th Dose)
Measles, Mumps,
Rubella (MMR)
Varicella (VAR)
Hepatitis A (HepA)



DTaP (4th Dose)



HepA (2nd Dose)
Total of 2 doses:
12-23 months



Influenza (Flu Shot) Yearly: 6 months to 18 years
Some kids under 8 may get 2 flu shots every year

Doctors may choose different types of vaccines that may have different dosages. So your child may or may not get an extra dose of some immunizations. Talk to your doctor for more information.

Source: Centers for Disease Control and Prevention (CDC)

300900THMDKYEN 221210



DTaP (5th Dose)
 IPV (4th Dose)
 MMR (2nd Dose)
 VAR (2nd Dose)

Tetanus, Diphtheria,
 Acellular Pertussis
 (Tdap)
 Meningococcal
 Human
 Papillomavirus
 (HPV) 2 doses



Most children will get their HPV shot at age 11. Some children with certain risk factors can get the HPV shot as early as age 9. Some children may need a 3rd dose. Talk to your doctor to find out what is best for your child.

Meningococcal



Influenza (Flu Shot) Yearly: 6 months to 18 years
Some kids under 8 may get 2 flu shots every year



MARK04782 APP_X/XX/2021