

Original Effective Date: 07/01/2018 Current Effective Date: 09/25/2024 Last P&T Approval/Version: 07/31/2024

Next Review Due By: 07/2025 Policy Number: C13398-A

# Nitisinone (Orfadin, Nityr)

**PRODUCTS AFFECTED** 

Nityr (nitisinone), nitisinone, Orfadin (nitisinone)

## **COVERAGE POLICY**

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Coverage Guideline must be read in its entirety to determine coverage eligibility, if any. This Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide Molina Healthcare complete medical rationale when requesting any exceptions to these guidelines.

## **Documentation Requirements:**

Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

#### **DIAGNOSIS:**

Hereditary tyrosinemia type 1

#### **REQUIRED MEDICAL INFORMATION:**

This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. If a drug within this policy receives an updated FDA label within the last 180 days, medical necessity for the member will be reviewed using the updated FDA label information along with state and federal requirements, benefit being administered and formulary preferencing. Coverage will be determined on a case-by case basis until the criteria can be updated through Molina Healthcare, Inc. clinical governance. Additional information may be required on a case-by-case basis to allow for adequate review. When the requested drug product for coverage is dosed by weight, body surface area or other member specific measurement, this data element is required as part of the medical necessity review. The Pharmacy and Therapeutics Committee has determined that the drug benefit shall be a mandatory generic and that generic drugs will be dispensed whenever available.

#### A. HEREDITARY TYROSINEMIA TYPE 1:

 Documented diagnosis of hereditary tyrosinemia type 1 (HT-1) AND

Molina Healthcare, Inc. confidential and proprietary © 2024

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

## Drug and Biologic Coverage Criteria

- Documentation diagnosis was confirmed by detection of elevated succinyl acetone (SA) in blood or urine OR DNA testing confirming mutation in the fumarylacetoacetate hydrolase (FAH) gene [DOCUMENTATION REQUIRED]
- Documentation of baseline succinyl acetone (SA) level [DOCUMENTATION REQUIRED]
   AND
- Prescriber attestation member has been counseled regarding dietary restriction of tyrosine and phenylalanine AND
- Prescriber attests baseline ophthalmologic testing, hepatic imaging and baseline labs have been obtained and reviewed such as: liver evaluation (PT, PTT, ALT/AST), renal function (BUN, creatine, etc.), plasma amino acids, and a complete blood count (CBC), and serum alpha- fetoprotein (AFP) AND
- 6. FOR ORFADIN REQUESTS: Clinical evidence or medical record documenting the use of Nityr will be ineffective or cause an adverse reaction to the member

## **CONTINUATION OF THERAPY:**

## A. HEREDITARY TYROSINEMIA TYPE 1:

- Prescriber attestation that there has been monitoring for plasma amino acids, liver function, serum AFP increases, CBC, and ophthalmologic side effects testing
  - Note: Patients with hereditary tyrosinemia type I are at increased risk of developing porphyric crises, hepatic neoplasms, and liver failure requiring liver transplantation. Regular monitoring of the liver by imaging and laboratory tests, including serum alpha-fetoprotein concentrations, is recommended. An increase in alpha-fetoprotein concentrations may be a sign of inadequate nitisinone treatment, but patients with increasing alpha-fetoprotein concentrations or signs of nodules in the liver during treatment with nitisinone should always be evaluated for hepatic malignancy.

**AND** 

- Adherence to therapy at least 85% of the time as verified by prescriber and member's medication fill history (review Rx history for compliance) AND
- Documentation urinary or blood succinyl acetone (SA) levels have decreased from baseline while on treatment with nitisinone [DOCUMENTATION REQUIRED]
   AND
- Prescriber attests to or clinical review has found no evidence of intolerable adverse effects or drug toxicity (e.g., corneal ulcers, corneal opacities, keratitis, conjunctivitis, ocular pain, photophobia, etc.)

## **DURATION OF APPROVAL:**

Initial authorization: 6 months, Continuation of therapy: 12 months

## PRESCRIBER REQUIREMENTS:

Prescribed by, or in consultation with, a specialist in metabolic or genetic disease, or in the treatment of hereditary tyrosinemia type 1 (HT-1) [If prescribed in consultation, consultation notes must be submitted with initial request and reauthorization requests]

#### **AGE RESTRICTIONS:**

No restriction

#### **QUANTITY:**

Maximum 1 mg/kg orally twice daily (2 mg/kg/day)

## PLACE OF ADMINISTRATION:

The recommendation is that oral medications in this policy will be for pharmacy benefit coverage and

Molina Healthcare, Inc. confidential and proprietary © 2024

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Drug and Biologic Coverage Criteria patient self-administered.

## **DRUG INFORMATION**

#### **ROUTE OF ADMINISTRATION:**

Oral

#### **DRUG CLASS:**

Hereditary Tyrosinemia Type 1 (HT-1) Treatment - Agents

#### FDA-APPROVED USES:

Indicated for the treatment of adult and pediatric patients with hereditary tyrosinemia type 1 in combination with dietary restriction of tyrosine and phenylalanine

## **COMPENDIAL APPROVED OFF-LABELED USES:**

None

## **APPENDIX**

#### **APPENDIX:**

None

## **BACKGROUND AND OTHER CONSIDERATIONS**

#### **BACKGROUND:**

Hereditary Tyrosinemia Type 1 (HT-1): a rare metabolic disease in children. In normal, unaffected individuals, excess amounts of the amino acid tyrosine are degraded in several steps. In HT-1, however, one of the enzymes in this degradation, fumarylacetoacetase hydrolase (FAH), is deficient. Tyrosine and its toxic metabolites [fumarylacetoacetate, maleylacetoacetate, succinyl acetone (SA), and succinyl acetoacetate (SAA)] thus build up in the body and cause serious medical problems such as liver failure and hepatocellular carcinoma. Kidney dysfunction, skeletal changes, and neurological manifestations may also occur. Orfadin and Nityr are both indicated for the treatment of hereditary tyrosinemia type 1 (HT-1) in combination with dietary restriction of tyrosine and phenylalanine. HT-1 is the most severe disorder of tyrosine metabolism. Fumarylacetoacetate (FAA) causes damage as it accumulates in the liver and kidney. FAA also causes oxidative damage to cells. To diagnose a patient with HT-1 the metabolites ofFAA, succinyl acetoacetate (SAA) and succinyl acetone (SA), can be measured.

Nitisinone is the primary treatment for HT-1 as it limits formation of the toxic compounds such as FAA and its metabolite SA.

# CONTRAINDICATIONS/EXCLUSIONS/DISCONTINUATION:

All other uses of Nitisinone (Orfadin, Nityr) are considered experimental/investigational and therefore, will follow Molina's Off- Label policy. Contraindications to Nitisinone (Orfadin, Nityr) include: No labeled contraindications.

#### OTHER SPECIAL CONSIDERATIONS:

None

#### **CODING/BILLING INFORMATION**

Note: 1) This list of codes may not be all-inclusive. 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement

HCPCS CODE	DESCRIPTION
NA	

# Drug and Biologic Coverage Criteria

## **AVAILABLE DOSAGE FORMS:**

Nityr TABS 2MG, 5MG & 10MG Nitisinone CAPS 2MG, 5MG, 10MG, & 20MG Orfadin CAPS 2MG, 5MG, 10MG, & 20MG Orfadin SUSP 4MG/ML

## **REFERENCES**

- 1. Nityr [packaged insert]. Cycle Pharmaceuticals Ltd. Cambridge, UK. January 2024.
- 2. Orfadin [package insert]. Sobi, Inc. Waltham, MA. November 2021.
- 3. Chinsky, JM et al. Diagnosis and treatment of tyrosinemia type I: a US and Canadian consensus group review and recommendations. Genet Med. 2017 Dec; 19(12). doi: [10.1038/gim.2017.101]

SUMMARY OF REVIEW/REVISIONS	DATE
REVISION- Notable revisions:	Q3 2024
Required Medical Information	
References	
REVISION- Notable revisions:	Q3 2023
Products Affected	
Required Medical Information	
Continuation of Therapy	
FDA-Approved Uses	
Available Dosage Forms	
References	
REVISION- Notable revisions:	Q3 2022
Required Medical Information	
Continuation of Therapy	
Quantity	
Contraindications/Exclusions/Discontinuation	
References	
Q2 2022 Established tracking in new	Historical changes on file
format	